

12-DIFFERENTIATION

Differentiation and Applications of Derivatives :

- If $y = f(x)$, then

$$1. \frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$2. \left(\frac{dy}{dx} \right)_{x=a} = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

$$3. \left(\frac{dy}{dx} \right)_{x=a} = \lim_{x \rightarrow h} \frac{f(a+h) - f(a)}{h}$$

- If $u = f(x)$, $v = \phi(x)$, then

$$1. \frac{d}{dx}(k) = 0$$

$$2. \frac{d}{dx}(ku) = k \frac{du}{dx}$$

$$3. \frac{d}{dx}(u \pm v) = \frac{du}{dx} \pm \frac{dv}{dx}$$

$$4. \frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

$$5. \frac{du}{dx} \left(\frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

$$6. \text{ If } x = f(t), y = \phi(t), \text{ then } \frac{dy}{dx} = \frac{dy}{dt} \bigg/ \frac{dx}{dt}$$

$$7. \text{ If } y = f[\phi(x)], \text{ then } \frac{dy}{dx} = f'[\phi(x)] \cdot \frac{d}{dx}[\phi(x)]$$

$$8. \text{ If } w = f(y), \text{ then } \frac{dw}{dx} = f'(y) \frac{dy}{dx}$$

$$9. \text{ If } y = f(x), z = \phi(x), \text{ then } \frac{dy}{dz} = \frac{dy}{dx} \cdot \frac{dx}{dz}$$

$$10. \frac{dy}{dx} \cdot \frac{dx}{dy} = 1 \text{ or } \frac{dy}{dx} = \frac{1}{dx/dy}$$

- 1. $\frac{d}{dx}(k) = 0$

$$2. \frac{d}{dx} x^n = nx^{n-1}$$

$$3. \frac{d}{dx} \frac{1}{x^n} = -\frac{n}{x^{n+1}}$$

$$4. \frac{d}{dx} \sqrt{x} = \frac{1}{2\sqrt{x}}$$

$$5. \frac{d}{dx} e^x = e^x$$

$$6. \frac{d}{dx} a^x = a^x \log a$$

$$7. \frac{d}{dx} \log x = \frac{1}{x}$$

$$8. \frac{d}{dx} \log_a x = \frac{1}{x} \log_a e$$

$$9. \frac{d}{dx} \sin x = \cos x$$

$$10. \frac{d}{dx} \cos x = -\sin x$$

$$11. \frac{d}{dx} \tan x = \sec^2 x$$

$$12. \frac{d}{dx} \cot x = -\operatorname{cosec}^2 x$$

$$13. \frac{d}{dx} \sec x = \sec x \tan x$$

$$14. \frac{d}{dx} \operatorname{cosec} x = -\operatorname{cosec} x \cot x$$

$$15. \frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}$$

$$16. \frac{d}{dx} \cos^{-1} x = -\frac{1}{\sqrt{1-x^2}}$$

$$17. \frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2}$$

$$18. \frac{d}{dx} \cot^{-1} x = -\frac{1}{1+x^2}$$

$$19. \frac{d}{dx} \sec^{-1} x = \frac{1}{x\sqrt{x^2-1}}$$

$$20. \frac{d}{dx} \operatorname{cosec}^{-1} x = -\frac{1}{x\sqrt{x^2-1}}$$

- **Suitable substitutions** : The functions any also be reduced to simpler forms by the substitutions as follows.

1. If the function involve the term $\sqrt{(a^2 - x^2)}$, then put $x = a \sin \theta$ or $x = a \cos \theta$.

2. If the function involve the term $\sqrt{(a^2 + x^2)}$, then put $x = a \tan \theta$ or $x = a \cot \theta$.

3. If the function involve the term $\sqrt{(x^2 - a^2)}$, then put $x = a \sec \theta$ or $x = a \operatorname{cosec} \theta$.

4. If the function involve the term $\sqrt{\frac{a-x}{a+x}}$, then put $x = a \cos \theta$ or $x = a \cos 2\theta$

All the above substitutions are also true, if $a = 1$

- **Differentiation by taking logarithm** :

Differentiation of the functions of the following types are obtained by taking logarithm.

1. When the functions consists of the product and quotient of a number of functions.
2. When a function of x is raised to a power which is itself a function of x .

For example, let $y = [f(x)]^{\phi(x)}$

Taking logarithm of both sides, $\log y = \phi(x) \log f(x)$

Differentiating both sides w.r.t 'x',

$$\frac{1}{y} \frac{dy}{dx} = \phi'(x) \log f(x) + \phi(x) \cdot \frac{f'(x)}{f(x)}$$

$$= [f(x)]^{\phi(x)} \log f(x) \cdot \phi'(x) + \phi(x) \cdot [f(x)]^{\phi(x)-1} \cdot f'(x)$$

$$\frac{dy}{dx} = \text{Differential of } y \text{ treating } f(x) \text{ as constant} +$$

Differential of y treating $\phi(x)$ as constant.

It is an important formula.

- **Differentiation of implicit functions** :

1. If $f(x, y) = 0$ is a implicit function, then

$$\frac{dy}{dx} = - \frac{\partial f / \partial x}{\partial f / \partial y}$$

$$= - \frac{\text{Diff. of } f \text{ w.r.t. } x \text{ keeping } y \text{ constant}}{\text{Diff. of } f \text{ w.r.t. } y \text{ keeping } x \text{ constant}}$$

For example, consider $f(x, y) = x^2 + 3xy + y^2 = 0$, then

$$\frac{dy}{dx} = - \frac{\partial f / \partial x}{\partial f / \partial y} = - \frac{2x + 3y}{3x + 2y}$$

1. If $y = f(x)$, then

$$\frac{dy}{dx} = y_1 = f'(x), \quad \frac{d^2y}{dx^2} = y_2 = f''(x), \dots$$

$$\frac{d^2y}{dx^2} = y_2 = f''(x)$$

$$2. \frac{d^n}{dx^n} (ax + b)^n = n! a^n$$

$$3. \frac{d^n}{dx^n} (ax + b)^m = m(m-1) \dots (m-n+1) a^n (ax + b)^{m-n}$$

$$4. \frac{d^n}{dx^n} e^{mx} = m^n e^{mx}$$

$$5. \frac{d^n}{dx^n} a^{mx} = m^n a^{mx} (\log a)^n$$

$$6. \frac{d^n}{dx^n} \log(ax + b) = \frac{(-1)^{n-1} a^n (n-1)!}{(ax + b)^n}$$

$$7. \frac{d^n}{dx^n} \sin(ax + b) = a^n \sin\left(ax + b + \frac{n\pi}{2}\right)$$

$$8. \frac{d^n}{dx^n} \cos(ax + b) = a^n \cos\left(ax + b + \frac{n\pi}{2}\right)$$

- **Leibnitz's theorem** : If u and v are any two functions of x such that their desired differential coefficients exist, then the n^{th} differential coefficient of uv is given by

$$D^n(uv) = (D^n u)v + {}^n C_1 (D^{n-1} u)(Dv) + {}^n C_2 (D^{n-2} u)(D^2 v) + \dots + u(D^n v)$$

Do you know



- Did you know that there are 206 bones in the adult human body and there are 300 in children (as they grow some of the bones fuse together).
- Flea's can jump 130 times higher than their own height. In human terms this is equal to a 6ft. person jumping 780 ft. into the air.
- The most dangerous animal in the world is the common housefly. Because of their habits of visiting animal waste, they transmit more diseases than any other animal.
- Snakes are true carnivorous because they eat nothing but other animals. They do not eat any type of plant material.
- The world's largest amphibian is the giant salamander. It can grow up to 5 ft. in length.
- The smallest bone in the human body is the stapes or stirrup bone located in the middle ear. It is approximately .11 inches (.28 cm) long.